

Appendices index

Appendix 1 Search strategies.....	2
Appendix 2 Characteristics of included studies	5
Appendix 3 List of included studies.....	15
Appendix 4 List of excluded studies.....	20
Appendix 5 Network plots of eligible comparisons for five outcomes.....	24
Appendix 6 Risk of bias assessment	27
Appendix 7 Pairwise meta-analysis results for direct comparisons of interventions.....	29
Appendix 8 Ranking of treatments for pregnancy	31
Appendix 9 Network plot for pregnancy incorporating risk of bias assessment	32
Appendix 10 Inconsistency plot for pregnancy.....	33
Appendix 11 Comparison-adjusted funnel plot for pregnancy.....	34
Appendix 12 Network meta-analysis results for live birth.....	35
Appendix 13 Ranking of treatments for live birth	36
Appendix 14 Network meta-analysis results for ovulation.....	37
Appendix 15 Ranking of treatments for ovulation	38
Appendix 16 Network meta-analysis results for miscarriage per woman randomised.....	39
Appendix 17 Network meta-analysis results for miscarriage per pregnancy.....	40
Appendix 18 Network meta-analysis results for multiple pregnancy.....	41
Appendix 19 Ranking of treatments for multiple pregnancy	42
Appendix 20 Network plot for live birth incorporating risk of bias assessment	43
Appendix 21 Inconsistency plot for live birth.....	44
Appendix 22 Comparison-adjusted funnel plot for live birth.....	45
Appendix 23 Network plot for ovulation incorporating risk of bias assessment	46
Appendix 24 Inconsistency plot for ovulation.....	47
Appendix 25 Comparison-adjusted funnel plot for ovulation.....	48
Appendix 26 Network plot for miscarriage incorporating risk of bias assessment	49
Appendix 27 Ranking of treatments for miscarriage per pregnancy	50
Appendix 28 Inconsistency plot for miscarriage per pregnancy	51
Appendix 29 Comparison-adjusted funnel plot for miscarriage per pregnancy	52
Appendix 30 Network plot for multiple pregnancy incorporating risk of bias assessment	53
Appendix 31 Inconsistency plot for multiple pregnancy	54
Appendix 32 Comparison-adjusted funnel plot for multiple pregnancy	55
Appendix 33 Sensitivity analysis - RCTs reporting clinical pregnancy.....	56
Appendix 34 Sensitivity analysis - RCTs with treatment naïve women	57
Appendix 35 Sensitivity analysis - RCTs with low risk of randomisation & allocation bias....	58
Appendix 36 Additional discussion.....	59
Appendix 37 Meta-analysis of clomiphene and metformin combined vs clomiphene alone for discontinuation due to side effects.	61
Appendix 38 Side effects of clomiphene and metformin combined vs clomiphene alone.....	62
Appendix 39 Congenital malformations in newborns conceived through letrozole vs control..	64

Appendix 1 Search strategies

1a. MEDLINE search strategy Database: Ovid MEDLINE(R)

- 1 exp Polycystic Ovary Syndrome/
- 2 Polycystic Ovar\$.tw.
- 3 PCOS.tw.
- 4 PCOD.tw.
- 5 PCO.tw.
- 6 (stein-leventhal or leventhal).tw.
- 7 (ovar\$ adj (scelerocystic or polycystic or degeneration)).tw.
- 8 anovulat\$.ti,ab,sh,tw.
- 9 oligo ovulat\$.ti,ab,sh,tw.
- 10 or/1-9
- 11 randomized controlled trial.pt.
- 12 controlled clinical trial.pt.
- 13 randomly.ab,ti.
- 14 randomized.ab,ti.
- 15 (crossover or cross over).tw.
- 16 placebo.tw.
- 17 RCT.tw.
- 18 trial.ti.
- 19 clinical trials as topic.sh.
- 20 or/11-19
- 21 exp animals/ not humans.sh.
- 22 20 not 21
- 23 fertil\$.ti,ab,sh,tw.
- 24 infertil\$.ti,ab,sh,tw.
- 25 subfertil\$.ti,ab,sh,tw.
- 26 pregnan\$.ti,ab,sh,tw.
- 27 exp ovulation induction/ or exp superovulation/
- 28 (ovulat\$ adj2 induc\$).tw.
- 29 (ovar\$ adj2 stimulat\$).tw.
- 30 superovulat\$.tw.
- 31 or/23-30
- 32 10 and 22 and 31

1b. Embase search strategy Database: EMBASE.com

#1 'ovary polycystic disease'/exp OR 'stein leventhal syndrome'/exp
 #2 (polycystic NEAR/2 ovar*):de,ab,ti
 #3 pcod:de,ab,ti OR pcod:de,ab,ti OR pco:de,ab,ti
 #4 leventhal:de,ab,ti
 #5 (ovar* NEAR/2 (scelerocystic OR degeneration)):de,ab,ti
 #6 'anovulation'/exp
 #7 anovulat*:de,ab,ti
 #8 (oligo NEAR/2 ovulat*):de,ab,ti
 #9 #1 OR #2 OR #3 OR #4 OR #5 OR #6 OR #7 OR #8
 #10 'randomized controlled trial'/exp
 #11 'controlled clinical trial'/exp
 #12 randomized:de,ab,ti
 #13 randomly:de,ab,ti
 #14 trial:ti
 #15 placebo:de,ab,ti
 #16 rct:de,ab,ti
 #17 crossover:de,ab,ti OR (cross NEAR/1 over):de,ab,ti
 #18 'clinical trial' OR 'clinical trials':de
 #19 #10 OR #11 OR #12 OR #13 OR #14 OR #15 OR #16 OR #17 OR #18
 #20 #19 AND [animals]/lim NOT [humans]/lim
 #21 #19 NOT #20
 #22 'infertility'/exp OR 'fertility'/exp OR 'subfertility'/exp
 #23 infertil*:de,ab,ti OR subfertil*:de,ab,ti OR fertil*:de,ab,ti
 #24 pregnan*:de,ab,ti
 #25 'pregnancy'/exp
 #26 'ovulation induction'/exp OR 'superovulation'/exp
 #27 (ovulat* NEAR/2 induc*):de,ab,ti
 #28 (ovar* NEAR/2 stimulat*):de,ab,ti
 #29 superovulat*:de,ab,ti
 #30 #22 OR #23 OR #24 OR #25 OR #26 OR #27 OR #28 OR #29
 #31 #9 AND #21 AND #30

1c. Database: EBM Reviews - Cochrane Central Register of Controlled Trials

#1 [mh "Polycystic Ovary Syndrome"]
 #2 (polycystic near ovar*):kw,ab,ti
 #3 pcod:kw,ab,ti or pcod:kw,ab,ti or pco:kw,ab,ti
 #4 leventhal:kw,ab,ti
 #5 (ovar* near (scelerocystic or degeneration)):kw,ab,ti
 #6 anovulat*:kw,ab,ti
 #7 oligo near ovulat*:kw,ab,ti
 #8 [mh anovulation]
 #9 #1 or #2 or #3 or #4 or #5 or #6 or #7 or #8
 #10 randomized controlled trial:pt
 #11 controlled clinical trial:pt

#12 placebo:kw,ti,ab
#13 randomly:kw,ti,ab
#14 RCT:kw,ti,ab
#15 trial:ti
#16 crossover:kw,ti,ab or (cross next over):kw,ti,ab
#17 #10 or #11 or #12 or #13 or #14 or #15 or #16
#18 [mh infertility]
#19 [mh fertility]
#20 [mh pregnancy]
#21 infertil*:kw,ti,ab
#22 fertil*:kw,ti,ab
#23 subfertil*:kw,ti,ab
#24 pregnan*:kw,ti,ab
#25 [mh "Ovulation Induction"] or [mh superovulation]
#26 ovulat* near induc*:kw,ti,ab
#27 ovar* near stimulat*:kw,ti,ab
#28 superovulat*:kw,ti,ab
#29 #18 or #19 or #20 or #21 or #22 or #23 or #24 or #25 or #26 or #27 or #28
#30 #9 and #17 and #29

Appendix 2 Characteristics of included studies

Study	Interventions	Age (mean)	BMI (mean)	DOI (mean years)	Inclusion criteria	Sample Size	Previous Treatment	Country	Setting	Maximum of treatment cycles	IUI or TI
Abuelghar 2013 ¹	CC MF+CC	28.4 27.6	28.1 28.6	2.8 3.2	Overweight and obese infertile women with PCOS (Rotterdam criteria)	66	unknown	Egypt	single-centre	1	TI
Amer 2009 ²	CC LOD	29.1 28.1	26.1 26.2	1.8 2.1	PCOS (at least 2 of the following 3 features: clinical [oligo/amenorrhoea and/or Hyperandrogenaemia], biochemical [LH≥10 IU/l, LH/FSH ratio ≥2, testosterone>2.6 nmol/l or free androgen index (FAI) >5] and/or sonographic (polycystic ovaries) features.)	72	naive	UK	single-centre	6	TI
Amer 2015 ³	CC LET	NA	NA	NA	anovulatory women with PCOS	159	naive	UK	single-centre	7	TI
Atay 2006 ⁴	CC LET	26.2 27.1	25.8 26.1	2.4 2.2	Women with primary infertility and PCOS(oligo- or amenorrhoea and ovaries with at least 10 subcapsular cysts 2 – 10 mm in diameter	106	unknown	Turkey	N/A	1	TI

					and hyperechogenic stroma.)						
Ayaz 2013 ⁵	CC MF+CC	31.3 32.0	NA ^a	NA	PCOS (the presence of two of the three following criteria:1. Polycystic ovaries [either 12 or more peripheral follicles or increased ovarian volume, > 10 cm ³]. 2. Oligo or anovulation [irregular cycles, amenorrhea]. 3. Clinical and/or biochemical signs of hyperandrogenism [Acne, hirsutism, voice changes, and Clitoromegaly].)	42	unknown	Saudi Arabia	single-centre	6	TI
Aygen 2007 ⁶	CC LET	23.4 26.8	27.6 26.9	4.2 5.8	Infertility and PCOS (Rotterdam criteria)	10	unknown	Turkey	single-centre	6	TI
Badawy 2009 ⁷	CC LET	29.3 27.1	27.1 28.1	NA	Infertile women with PCOS (Rotterdam criteria)	438	unknown	Egypt	multi-centre	>1	TI
Badawy 2011 ⁸	CC TAM	25.8 26.2	29.9 30.5	1.5 1.4	PCOS (Rotterdam criteria)	371	unknown	Egypt	multi-centre	1	TI
Basirat 2012 ⁹	CC MF+CC	25.3 24.9	25.4 26.3	2.7 2.4	Infertile PCOS (Rotterdam criteria)	334	unknown	Iran	multi-centre	3	IUI
Bayar 2006 ¹⁰	CC LET	30.6 32.2	NA	3 5	anovulatory PCOS (Rotterdam criteria)	80	naive	Turkey	single-centre	>1	TI
Beigi 2006 ¹¹	CC MF	NA	NA	NA	PCOS based on a history of hyperandrogenism, anovulation, oligomenorrhea	70	unknown	Iran	single-centre	6	TI

					or amenorrhea, diagnostic ultrasound and laboratory findings						
Boonstanfar 2001 ¹²	CC TAM	26.5 26.6	30.2 30.9	3.7 3.5	anovulatory women with infertility	95	naive	USA	single-centre	>1	TI
Boudhraa 2010 ¹³	CC MF+CC	30.7 30.6	29.8 30.0	2.5 ^b	PCOS (Rotterdam criteria) with subfertility	63	unknown	Tunis	single-centre	3-6	TI
Cudmore 1966 ¹⁴	CC PB	24.6 24.6	NA	NA	A diagnosis of secondary amenorrhea of at least 2 year's duration; persistent oligomenorrhea with no more than 4 periods in 1 year; or anovulatory infertility (infertility of more than 2 years' duration in which anovulation was the only cause found)	22	unknown	Canada	single-centre	3	TI
Dasari 2009 ¹⁵	CC MF+CC	NA ^c	NA ^d	NA	Infertile PCOS (Rotterdam criteria)	40	unknown	India	single-centre	6	TI
Dehbashi 2009 ¹⁶	CC LET	24.3 23.6	27.1 27.5	2.3 2.0	PCOS (Rotterdam criteria)	100	naive	Iran	single-centre	1	TI
El-Biely 2001 ¹⁷	CC MF+CC	25.7 26.4	27.4 28.7	4.7 4.5	Infertile obese patients with PCOS (oligomenorrhoea, ultrasound findings of ≥ 10 ovarian cysts measuring 2-8mm around a dense stroma)	90	unknown	Egypt	single-centre	6	TI

Fleming 2002 ¹⁸	MF	28.6	34.2	NA	Women with oligomenorrhea or amenorrhea and PCO	42	naive	UK	single-centre	4	TI
	PB	29.2	35.0								
Garcia 1985 ¹⁹	CC	27.6 ^e	NA	NA	Anovulatory infertile women	49	unknown	USA	single-centre	5	TI
	PB										
Homburg 2012 ²⁰	CC	29.4	25.7	2.1	anovulatory or oligo-ovulatory infertile women with PCOS (Rotterdam criteria)	302	naive	Netherlands, UK, Malta, Belgium, Argentina, Colombia	multi-centre	3	TI/IU
	FSH	29.8	25.1	2.1							
Jahan 2015 ²¹	CC	NA	NA	NA	PCOS	460	naive	Bangladesh	single-centre	6	TI
	LET MF										
Johnson 1966 ²²	CC	NA	NA	NA	Anovulatory women	65	mixed	USA	single-centre	1	TI
	PB										
Johnson 2010A ²³	MF	29.5	38.0	3.3(2.4	anovulatory or oligo-ovulatory women with PCOS (Rotterdam criteria), BMI>32 kg/m ²	65	mixed	New Zealand	multi-centre	6	TI
	PB	29.2	37.6	-5.9) ^f 3.4(2-5) ^f							
Johnson 2010B ²³	CC	28.2	26.2	2(1-3) ^f	anovulatory or oligo-ovulatory women with PCOS (Rotterdam criteria), BMI≤32 kg/m ²	106	mixed	New Zealand	multi-centre	6	TI
	MF	28.9	26.5	1.5(1-							
	MF+CC	29.2	26.9	4) ^f 2(1.5-5) ^f							
Kar 2012 ²⁴	CC	26.3	26.0	3.1	infertile PCOS (Rotterdam criteria)	103	naive	India	single-centre	1	TI/IU
	LET	26.3	25.9	3.1							

Kar 2015 ²⁵	CC	25.8	26.5	2.8	PCOS (Rotterdam criteria), with the primary complaints of infertility and oligomenorrhea	105	naive	India	single- centre	6	TI
	MF	25.2	24.5	1.7							
	MF+CC	26.6	27.2	2.5							
Karimzadeh 2007 ²⁶	MF	27.2	28.8	5.6	PCOS (Rotterdam criteria)	200	unknown	Iran	single- centre	3	TI
	PB	28.6	29.5	6.2							
Karimzadeh 2010 ²⁷	CC	27.5	27.2	4.1	infertile PCOS (Rotterdam criteria)	268	unknown	Iran	single- centre	6	TI
	MF	27.3	27.2	3.9							
	MF+CC	27.3	28.0	4.6							
Keikha 2011 ²⁸	CC	27.1	NA	2.9	infertile PCOS	116	naive	Iran	single- centre	1	TI
	LET	27.6		3.0							
Khorram 2006 ²⁹	CC	28.0	38.8	NA	PCOS (anovulatory or oligo- ovulatory cycles, polycystic ovaries on a baseline ultrasound, hyperandrogenism) and infertility	31	naive	USA	single- centre	1	TI
	MF+CC	28.4	35.3								
Leanza 2014 ³⁰	CC	26-34 ^g	NA	NA	PCOS (typical ultrasound situation, oligomenorrhea/amenorrhea, hyperandrogenism) with above 3 years of infertility, BMI>27.5	56	naive	Italy	single- centre	3	IUI
	MF+CC										
Legro 2007 ³¹	CC	27.9	36.0	3.5	infertile women PCOS (oligomenorrhea and hyperandrogenemia)	626	mixed	USA	multi- centre	6	TI
	MF	28.1	35.6	3.3							
	MF+CC	28.3	34.2	3.4							

Legro 2014 ³²	CC LET	28.8 28.9	35.1 35.2	3.5 3.4	infertile women PCOS (Rotterdam criteria)	750	mixed	USA	multi- centre	5	TI
Liu 2015 ³³	CC LET	NA	NA	NA	PCOS patients who have conception desire	134	unknown	China	single- centre	>1	TI
López 2004 ³⁴	CC FSH	29(23- 38) ^f 30(22- 39) ^f	22.3 21.9	3(1-8) ^f 3(1-8) ^f	anovulatory infertility due to PCOS (Rotterdam criteria)	76	naive	Spain	single- centre	3	TI
Lord 2006 ³⁵	MF PB	27.8 30.6	33.7 36.4	NA	PCOS (anovulation and a raised free androgen index (FAI) >5.0)	44	unknown	UK	single- centre	3	TI
Lorzadeh 2011 ³⁶	CC LET	26.1 28.2	25.4 24.2	NA	PCOS (based on the chronic anovulation and clinical/lab- based hyperandrogenism), age <35, No successful pregnancy after one year of weekly (2-3 times) sexual contact without contraception.	100	unknown	Iran	single- centre	>1	TI
Maged 2015 ³⁷	CC MF+CC	26.0 25.8	27.3 27.7	2.8 2.8	PCOS (Rotterdam criteria)	80	unknown	Egypt	single- centre	3	TI
Mobusher 2014 ³⁸	CC LET	24.3 24.3	25.9 25.9	3.1 3.2	PCOS (Rotterdam criteria) and infertility	100	naive	Pakistan	single- centre	1	TI
Moll 2006 ³⁹	CC MF+CC	28.4 27.9	27.8 28.5	1.3 1.6	PCOS (Rotterdam criteria), all women with chronic anovulation and polycystic	225	naive	Netherlan ds	multi- centre	6	TI

ovaries diagnosed by transvaginal ultrasonography											
Nazik 2012 ⁴⁰	CC	27.8	25.9	4.4	PCOS (Rotterdam criteria)	64	naive	Turkey	single-centre	>1	TI
	LET	25.6	24.7	3.4							
Palomba 2005 ⁴¹	CC	25.9	26.7	1.7	primary infertile anovulatory women with PCOS (NIH criteria)	100	naive	Italy	single-centre	6	TI
	MF	26.4	27.0	1.6							
Raja 2005 ⁴²	CC	26.9	NA	4.9	Infertility and PCOS (the presence of polycystic ovaries on ultrasonography with two or more of the following criteria: Oligomenorrhoea [<6 cycles in preceding year]; hirsutism; hyperandrogenism; Elevated LH or LH: FSH >2])	100	unknown	Pakistan	single-centre	6	TI
	MF+CC	26.5		4.2							
Ray 2012 ⁴³	CC	29(20-35) ^f	28.5(24.2-33.6) ^f	2.4	Infertile PCOS (Rotterdam criteria)	147	unknown	India	single-centre	>1	TI
	LET	28(19-35) ^f	28.8(23.2-34.6) ^f	2.2							
Robinson 2003 ⁴⁴	CC MF+CC	NA	NA	NA	Women with a one-year history of infertility and diagnosed with hyperandrogenic oligoovulatory or anovulatory cycles as the sole etiology for their	48	unknown	USA	single-centre	6	TI

					infertility						
Roy 2012 ⁴⁵	CC	26.5	25.4	5.8	infertility and anovulatory	212	unknown	India	single-	3	TI
	LET	26.1	25.8	6.4	PCOS (Rotterdam criteria), BMI<28				centre		
Sahin 2004 ⁴⁶	CC	24.5(19	25.7(23.	3.5(1-	Primary infertility and PCOS	21	unknown	Turkey	single-	6	TI
	MF+CC	-28) ^f	1-35.7) ^f	8) ^f	(on the basis of three or more				centre		
		27(21-	30.4(24.	5(2-	of the following criteria:						
		31) ^f	6-33.9) ^f	10) ^f	polycystic ovaries on pelvic						
					ultrasound examination,						
					oligo/amenorrhoea,						
					hirsutism,						
					hyperandrogenaemia (total						
					testosterone > 80 ng/dl						
					and/or free testosterone >						
					3.18 pg/ml)) and elevated						
					serum LH:FSH ratio (LH:FSH >						
					2))						
Santonocito 2009 ⁴⁷	CC	27.4	27.1	1.7	infertility and anovulatory	36	unknown	Italy	single-	6	TI
	MF	28.1	26.8	1.6	PCOS (Rotterdam criteria), BMI< 30 kg/m ²				centre		
Selim 2012 ⁴⁸	CC	25.1	23.8	2.6	Infertile women with PCOS	220	naive	Egypt	single-	1	TI
	LET	26.0	24.4	2.9	(Rotterdam criteria)				centre		
Seyedoshohadaei 2012 ⁴⁹	CC	24.7	NA	3.0	non-PCOS anovulatory	150	unknown	Iran	single-	6	TI
	LET	26.9		4.1	infertility, and ovary without				centre		
	TAM	25.4		3.0	evidence of polycystic ovaries						
Sharief 2015 ⁵⁰	CC	25.3	27.8	2.3	primary infertility and	75	unknown	Iraq	single-	6	TI

	LET	26.1	28.1	2.4	anovulation due to PCOS (ultrasonographic polycystic ovaries plus one or more of the following: oligomenorrhoea, positive progesterone, withdrawal bleeding, hirsutism/acne, obesity, and Luteinizing hormone/Follicle-stimulating hormone (LH/FSH) ratio >2 or raised circulating androgen, normal thyroid stimulating hormone)				centre		
Sh-El-Arab Elsedeek 2011 ⁵¹	CC LET	25.0 25.0	29.2 27.7	NA	Nulliparous PCOS (Rotterdam criteria), BMI ≤35	124	unknown	Egypt	single- centre	1	TI
Tang 2006 ⁵²	MF PB	29.7 29.8	37.6 38.9	4.5 4.9	anovulatory PCOS (polycystic ovaries on transvaginal scan, together with either oligomenorrhoea or amenorrhoea) and a BMI of >30,	143	naive	UK	multi- centre	6	TI
Vegetti 1999 ⁵³	CC TAM	NA	NA	NA	Infertility and normogonadotropic anovulation	95	naive	Italy	single- centre	>1	TI
Williams 2009 ⁵⁴	CC	NA	NA	NA	women with PCOS who are	55	unknown	USA	N/A	6	TI

	MF+CC	attempting to conceive.									
Zain 2009 ⁵⁵	CC	29.6	32.9	2.9	PCOS (Rotterdam criteria)	124	naive	Malaysia	single-centre	6	TI
	MF	27.8	33.9	3.1							
	MF+CC	29.3	33.0	3.3							
Zeinalzadeh 2010 ⁵⁶	CC	23.1	NA	2.6	PCOS (based on ultrasonography finding, oligomenorrhea and an increased LH/FSH ratio (>3))	107	naive	Iran	single-centre	1	IUI
	LET	23.8		2.4							

(Abbreviations: CC, clomiphene citrate; PB, placebo or no treatment; LET, letrozole; MF, metformin; TAM, tamoxifen; FSH, follicle stimulating hormone; LOD, laparoscopic ovarian drilling; NA, not available; BMI, body mass index; DOI: Duration of infertility)

- The percentages of women with BMI>25 in CC and CC+MF group are 71.4% and 56.7%, respectively.
- The mean duration of infertility of all the participants (including both groups).
- The percentages of women with age >31, 26-30 and 20-25 years are 8.3%, 41.7%, 50% in CC group and 18.8%, 43.8% and 37.5% in CC+MF group.
- The percentages of women with BMI >25 and BMI < 25 are 37.5% and 62.5%, respectively.
- in treatment group only
- median (range)
- range

Appendix 3 List of included studies

1. Abuelghar WM, Elkady OS, Khamees AA. Clomiphene citrate alone, in combination with metformin or in combination with pioglitazone as first line therapy in induction of ovulation in infertile women with polycystic ovary syndrome, a randomized controlled trial. *Middle East Fertility Society Journal* 2013;**18**(3):135-41
2. Amer S, Fakis A, Smith J, et al. Double blind cross-over randomized controlled trial comparing letrozole versus clomiphene citrate for ovulation induction in women with polycystic ovarian syndrome. *Human Reproduction* 2015;**30**:i96
3. Amer SA, Li TC, Metwally M, et al. Randomized controlled trial comparing laparoscopic ovarian diathermy with clomiphene citrate as a first-line method of ovulation induction in women with polycystic ovary syndrome. *Human Reproduction* 2009;**24**(1):219-25
4. Atay V, Cam C, Muhcu M, et al. Comparison of letrozole and clomiphene citrate in women with polycystic ovaries undergoing ovarian stimulation. *Journal of international medical research* 2006;**34**(1):73-6
5. Ayaz A, Alwan Y, Farooq MU. Metformin-clomiphene citrate vs. clomiphene citrate alone: Polycystic ovarian syndrome. *Journal of human reproductive sciences* 2013;**6**(1):15-18
6. Aygen EM, Güzel Z, Özgün T, et al. The use of letrozole for ovulation induction in infertile women with polycystic ovarian syndrome. *Erciyes Tip Dergisi* 2007;**29**(3):195-200
7. Badawy A, Abdel Aal I, Abulatta M. Clomiphene citrate or letrozole for ovulation induction in women with polycystic ovarian syndrome: a prospective randomized trial. *Fertility & Sterility* 2009;**92**(3):849-52
8. Badawy A, Gibreal A. Clomiphene citrate versus tamoxifen for ovulation induction in women with PCOS: a prospective randomized trial. *European Journal of Obstetrics, Gynecology, & Reproductive Biology* 2011;**159**(1):151-4 doi: <http://dx.doi.org/10.1016/j.ejogrb.2011.07.015>.
9. Basirat Z, Kashifard M, Amiri MG. Enhanced ovarian follicular development by Metformin does not correlate with pregnancy rate: A randomized trial. *International Journal of Fertility and Sterility* 2012;**6**(1):31-36
10. Bayar U, Basaran M, Kiran S, et al. Use of an aromatase inhibitor in patients with polycystic ovary syndrome: a prospective randomized trial. *Fertility & Sterility* 2006;**86**(5):1447-51
11. Beigi A. Randomized trial comparing clomiphene citrate and metformin as the first-line treatment for ovulation induction in polycystic ovary syndrome. *Human reproduction (Oxford, England)* 2006;**21**(Suppl):i129
12. Boostanfar R, Jain JK, Mishell DR, Jr., et al. A prospective randomized trial comparing clomiphene citrate with tamoxifen citrate for ovulation induction. *Fertility & Sterility* 2001;**75**(5):1024-6
13. Boudhraa K, Jellouli MA, Amri M, et al. [Indication of metformin in the management of hormonal dysfunction secondary to polycystic ovarian syndrome: prospective comparative study of 63 cases]. *Tunisie medicale* 2010;**88**(5):335-40
14. Cudmore DW, Tupper WR. Induction of ovulation with clomiphene citrate. A double-blind study. *Fertil Steril* 1966;**17**(3):363-73

15. Dasari P, Pranahita GK. The efficacy of metformin and clomiphene citrate combination compared with clomiphene citrate alone for ovulation induction in infertile patients with PCOS. *Journal of human reproductive sciences* 2009;**2**(1):18-22
16. Dehbashi S, Dehbashi S, Kazerooni T, et al. Comparison of the effects of letrozole and clomiphene citrate on ovulation and pregnancy rate in patients with polycystic ovary syndrome. *Iranian Journal of Medical Sciences* 2009;**34**(1):23-28
17. El-Biely MM, Habba M. The use of metformin to augment the induction of ovulation in obese infertile patients with polycystic ovary syndrome. *Middle East Fertility Society Journal* 2001;**6**(1):43-49
18. Fleming R, Hopkinson ZE, Wallace AM, et al. Ovarian function and metabolic factors in women with oligomenorrhea treated with metformin in a randomized double blind placebo-controlled trial. *Journal of Clinical Endocrinology & Metabolism* 2002;**87**(2):569-74
19. Garcia CR, Freeman EW, Rickels K, et al. Behavioral and emotional factors and treatment responses in a study of anovulatory infertile women. *Fertility & Sterility* 1985;**44**(4):478-83
20. Homburg R, Hendriks ML, Konig TE, et al. Clomifene citrate or low-dose FSH for the first-line treatment of infertile women with anovulation associated with polycystic ovary syndrome: a prospective randomized multinational study. *Human Reproduction* 2012;**27**(2):468-73
21. Jahan S. Comparative study of efficacy among metformin, clomiphene citrate and aromatase inhibitor (letrozole) as the first-line medication for ovulation induction, achievement of pregnancy and live birth in Asian women with polycystic ovarian syndrome: A prospective trial. *International Journal of Gynecology and Obstetrics* 2015;**131**:E503
22. Johnson JE, Jr., Cohen MR, Goldfarb AF, et al. The efficacy of clomiphene citrate for induction of ovulation. A controlled study. *Int J Fertil* 1966;**11**(3):265-70
23. Johnson NP, Stewart AW, Falkiner J, et al. PCOSMIC: a multi-centre randomized trial in women with PolyCystic Ovary Syndrome evaluating Metformin for Infertility with Clomiphene. *Human Reproduction* 2010;**25**(7):1675-83
24. Kar S. Clomiphene citrate or letrozole as first-line ovulation induction drug in infertile PCOS women: A prospective randomized trial. *Journal of human reproductive sciences* 2012;**5**(3):262-65
25. Kar S, Sanchita S. Clomiphene citrate, metformin or a combination of both as the first line ovulation induction drug for Asian Indian women with polycystic ovarian syndrome: A randomized controlled trial. *Journal of Human Reproductive Sciences* 2015;**8**(4):197-201
26. Karimzadeh MA, Eftekhari M, Taheripana R, et al. The effect of administration of metformin on lipid profile changes and insulin resistance in patients with polycystic ovary syndrome. *Middle East Fertility Society Journal* 2007;**12**(3):174-78
27. Karimzadeh MA, Javedani M. An assessment of lifestyle modification versus medical treatment with clomiphene citrate, metformin, and clomiphene citrate-metformin in patients with polycystic ovary syndrome. *Fertility & Sterility* 2010;**94**(1):216-20 doi: 10.1016/j.fertnstert.2009.02.078.

28. Keikha F, Shahraki Mojahed B. Induction ovulation in polycystic ovary patient with clomiphene citrate and letrozole. *Iranian Journal of Reproductive Medicine* 2011;**9**:46
29. Khorram O, Helliwell JP, Katz S, et al. Two weeks of metformin improves clomiphene citrate-induced ovulation and metabolic profiles in women with polycystic ovary syndrome. *Fertility & Sterility* 2006;**85**(5):1448-51 doi: 10.1016/j.fertnstert.2005.10.042.
30. Leanza V, Coco L, Grasso F, et al. Ovulation induction with clomiphene citrate and metformin in women with polycystic ovary syndrome. *Minerva ginecologica* 2014;**66**(3):299-301
31. Legro RS, Barnhart HX, Schlaff WD, et al. Clomiphene, metformin, or both for infertility in the polycystic ovary syndrome. *New England journal of medicine* 2007;**356**(6):551-66
32. Legro RS, Brzyski RG, Diamond MP, et al. Letrozole versus clomiphene for infertility in the polycystic ovary syndrome. *N Engl J Med* 2014;**371**(2):119-29 doi: 10.1056/NEJMoa1313517.
33. Liu C, Feng G, Wang Q, et al. Comparison of ovulation induction protocol for women with polycystic ovarian syndrome: A prospective randomized trial. *International Journal of Gynecology and Obstetrics* 2015;**131**:E231-E32
34. Lopez E, Gunby J, Daya S, et al. Ovulation induction in women with polycystic ovary syndrome: randomized trial of clomiphene citrate versus low-dose recombinant FSH as first line therapy. *Reproductive Biomedicine Online* 2004;**9**:382-90
35. Lord J, Thomas R, Fox B, et al. The effect of metformin on fat distribution and the metabolic syndrome in women with polycystic ovary syndrome--a randomised, double-blind, placebo-controlled trial. *BJOG: An International Journal of Obstetrics & Gynaecology* 2006;**113**(7):817-24
36. Lorzadeh N, Kazemirad S, Mohammadi Z. Comparison of effects letrozole and clomiphene citrate for ovulation induction in women with polycystic ovary syndrome. *Iranian Journal of Obstetrics, Gynecology and Infertility* 2011;**14**(3):13-9.
37. Maged AM, Elsayah H, Abdelhafez A, et al. The adjuvant effect of metformin and N-acetylcysteine to clomiphene citrate in induction of ovulation in patients with Polycystic Ovary Syndrome. *Gynecological endocrinology* 2015
38. Mobusher I. Comparison of the efficacy of letrozole and clomiphene citrate for ovulation induction in infertile women with polycystic ovary syndrome. *Pakistan Journal of Medical and Health Sciences* 2014;**8**(4):905-08
39. Moll E, Bossuyt PM, Korevaar JC, et al. Effect of clomifene citrate plus metformin and clomifene citrate plus placebo on induction of ovulation in women with newly diagnosed polycystic ovary syndrome: randomised double blind clinical trial. *BMJ (Clinical research ed)* 2006;**332**(7556):1485 doi: 10.1136/bmj.38867.631551.55.
40. Nazik H, Kumtepe Y. Comparison of efficacy of letrozole and clomiphene citrate in ovulation induction for women with polycystic ovarian syndrome. *HealthMED* 2012;**6**(3):879-83
41. Palomba S, Orio F, Jr., Falbo A, et al. Prospective parallel randomized, double-blind, double-dummy controlled clinical trial comparing clomiphene citrate and metformin as the first-line treatment for ovulation induction in nonobese anovulatory women with polycystic ovary syndrome. *Journal of Clinical Endocrinology & Metabolism*

- 2005;**90**(7):4068-74 doi: 10.1210/jc.2005-0110.
42. Raja A, Hashmi SN, Sultana N, et al. Presentation of polycystic ovary syndrome and its management with clomiphene alone and in combination with metformin. *J Ayub Med Coll Abbottabad* 2005;**17**(2):50-3
 43. Ray PB, Ray A, Chakraborti PS. Comparison of efficacy of letrozole and clomiphene citrate in ovulation induction in Indian women with polycystic ovarian syndrome. *Archives of Gynecology & Obstetrics* 2012;**285**(3):873-7 doi: <http://dx.doi.org/10.1007/s00404-011-2091-7>.
 44. Robinson R, Swezey M, Propst A, et al. Metformin added to clomiphene citrate does not improve pregnancy rates in hyperandrogenic, chronic anovulatory women: A randomized trial. *Fertility and sterility* 2003;**80**(Suppl 3):S273-S74, Abstract no: P
 45. Roy K, Baruah J, Singla S, et al. A prospective randomized trial comparing the efficacy of Letrozole and Clomiphene citrate in induction of ovulation in polycystic ovarian syndrome. *Journal of human reproductive sciences* 2012;**5**(1):20-25
 46. Sahin Y, Yirmibes U, Kelestimur F, et al. The effects of metformin on insulin resistance, clomiphene-induced ovulation and pregnancy rates in women with polycystic ovary syndrome. *European Journal of Obstetrics, Gynecology, & Reproductive Biology* 2004;**113**(2):214-20
 47. Santonocito V, Rapisarda V, Abruzzo SRM, et al. Comparison between clomiphene citrate and metformin for induction of ovulatory cycles in infertile nonobese women with polycystic ovary syndrome. *Giornale Italiano di Ostetricia e Ginecologia* 2009;**31**(11-12):455-60
 48. Selim MF, Borg TF. Letrozole and clomiphene citrate effect on endometrial and subendometrial vascularity in treating infertility in women with polycystic ovary syndrome. 2012; **28**(6): 405-10.
 49. Seyedoshohadaei F, Zandvakily F, Shahgeibi S. Comparison of the effectiveness of clomiphene citrate, tamoxifen and letrozole in ovulation induction in infertility due to isolated unovulation. *Iranian Journal of Reproductive Medicine* 2012;**10**(6):531-36
 50. Sharief M, Nafee NR. Comparison of letrozole and clomiphene citrate in women with polycystic ovaries undergoing ovarian stimulation. *Journal of the Pakistan Medical Association* 2015;**65**(11):1149-52
 51. Sheikh-El-Arab Elsedek M, Elmaghraby HAH. Predictors and characteristics of letrozole induced ovulation in comparison with clomiphene induced ovulation in anovulatory PCOS women. *Middle East Fertility Society Journal* 2011;**16**(2):125-30
 52. Tang T, Glanville J, Hayden CJ, et al. Combined lifestyle modification and metformin in obese patients with polycystic ovary syndrome. A randomized, placebo-controlled, double-blind multicentre study. *Human Reproduction* 2006;**21**(1):80-89
 53. Vegetti W, Riccaboni A, Columbo M, et al. Randomized study of induction of ovulation by two different molecules with antioestrogenic effects, in patients with chronic anovulation disorders. 1999; **72**(3 Suppl 1).
 54. Williams CD, Pastore LM, Shelly WB, et al. A randomized, placebo-controlled study of the influence of instant-release metformin on response to clomiphene citrate and time to conception in polycystic ovary syndrome. *Fertility and sterility* 2009;**92**(3):S105
 55. Zain MM, Jamaluddin R, Ibrahim A, et al. Comparison of clomiphene citrate, metformin,

- or the combination of both for first-line ovulation induction, achievement of pregnancy, and live birth in Asian women with polycystic ovary syndrome: a randomized controlled trial. *Fertility & Sterility* 2009;**91**(2):514-21
56. Zeinalzadeh M, Basirat Z, Esmailpour M. Efficacy of letrozole in ovulation induction compared to that of clomiphene citrate in patients with polycystic ovarian syndrome. *Journal of reproductive medicine* 2010;**55**(1-2):36-40

Appendix 4 List of excluded studies

1. Aboul Enien WM, Barghash NA, Mohamed Ali FS. Clinical, ultrasonographic and endocrine predictors of ovarian response to clomiphene citrate in normogonadotropic anovulatory infertility. *Middle East Fertility Society Journal* 2004;**9**(3):242-50
2. Alamolhoda S, Mirabi P. Metformin and/or Clomiphene do not adversely affect liver or renal function in women with polycystic ovary syndrome. *Iranian Journal of Reproductive Medicine* 2013;**11**:38
3. Al-Dahhan F. The role of Metformin in induction of ovulation in obese infertile patients with polycystic ovary syndrome. *International Journal of Gynecology and Obstetrics* 2009;**107**(Journal Article):S636
4. Amer SAK, Gopalan V, Li TC, et al. Long term follow-up of patients with polycystic ovarian syndrome after laparoscopic ovarian drilling: Clinical outcome. *Human Reproduction* 2002;**17**(8):2035-42
5. Ayaz A, Alwan Y, Farooq MU. Efficacy of combined metformin-clomiphene citrate in comparison with clomiphene citrate alone in infertile women with polycystic ovarian syndrome (PCOS). *Journal of Medicine & Life* 2013;**6**(2):199-201
6. Baillargeon JP, Jakubowicz DJ, Luorno MJ, et al. Effects of metformin and rosiglitazone, alone and in combination, in nonobese women with polycystic ovary syndrome and normal indices of insulin sensitivity. *Fertil Steril* 2004;**82**(4):893-902 doi: 10.1016/j.fertnstert.2004.02.127.
7. Baruah J, Roy KK, Rahman SM, et al. Endometrial effects of letrozole and clomiphene citrate in women with polycystic ovary syndrome using spiral artery Doppler. *Archives of gynecology and obstetrics* 2009;**279**(3):311-4 doi: 10.1007/s00404-008-0714-4.
8. Basirat Z, Golsourkhtabar M, Kashifard M. Does metformin affect on pregnancy outcome in PCOs infertile women with different BMI? *Iranian Journal of Reproductive Medicine* 2012;**10**:76
9. Begum MR, Ferdous J, Begum A, et al. Comparison of efficacy of aromatase inhibitor and clomiphene citrate in induction of ovulation in polycystic ovarian syndrome. *Fertility and sterility* 2009;**92**(3):853-57 doi: 10.1016/j.fertnstert.2007.08.044 [doi].
10. Beigi A, Zarrinkoub F. Randomized controlled trial comparing clomiphene citrate and metformin as the first-line treatment for ovulation induction in women with polycystic ovary syndrome. XVIII FIGO World Congress of Gynecology and Obstetrics 2006;**3**:177
11. Ben Ayed B, Dammak dit Mlik S, Ben Arab H, et al. Metformin effects on clomifene-induced ovulation in the polycystic ovary syndrome. *Tunisie Medicale* 2009;**87**(1):43-9
12. Connaughton JF, Jr., Garcia CR, Wallach EE. Induction of ovulation with cisclophene and a placebo. *Obstetrics & Gynecology* 1974;**43**(5):697-701
13. Diamond MP, Kruger M, Santoro N, et al. Adverse impact of progestin exposure and endometrial shedding prior to ovulation induction on conception and live birth in women with polycystic ovary syndrome. *Reproductive Sciences* 2012;**19**(3):97A
14. El Bigawy AF, Fouda UMF, Wahab HAE. A randomized trial of letrozole versus clomiphene citrate in induction of ovulation in patients with polycystic ovary syndrome (PCOS). *Middle East Fertility Society Journal* 2008;**13**(1):52-56

15. Ghahiri A, Mamorian M. Comparative study of aromatase inhibitor (Letrozole) with clomiphene citrate as the first line treatment of patients with PCO. *Iranian Journal of Reproductive Medicine* 2010;**8**:84
16. Hashim HA, Bazeed M, Elaali IA. Minimal stimulation or clomiphene citrate as first-line therapy in women with polycystic ovary syndrome: A randomized controlled trial. *Gynecological endocrinology* 2012;**28**(2):87-90
17. Homburg R, Hendriks ML, Konig T, et al. Clomifene or low-dose FSH for the first-line treatment of anovulatory PCOS: A prospective randomised multinational study (COFFI). *Molecular Human Reproduction* 2009;**24**:i22-i23
18. Hosseini MA, Alleyassin A, Sarvi F, et al. Metformin treatment in different phenotypes of polycystic ovary syndrome. *Archives of Gynecology and Obstetrics* 2013;**288**(5):1131-36
19. Imani B, Eijkemans MJC, Te Velde ER, et al. A nomogram to predict the probability of live birth after clomiphene citrate induction of ovulation in normogonadotropic oligoamenorrheic infertility. *Fertility and sterility* 2002;**77**(1):91-97
20. Johnson N. PCOSMIC polycystic ovarian syndrome, metformin for infertility with clomiphene: a multi centre randomised controlled trial. FSA Fertility Society Australia, Abstract Book of Proceedings Brisbane, Australia 19-22 October 2008 2008:34
21. Johnson NP. PCOSMIC-polycystic ovarian syndrome, metformin for infertility with clomiphene: A multi-centre double-blind randomised controlled trial. *Molecular Human Reproduction* 2009;**24**:i24
22. Jungheim ES, Odibo AO. Fertility treatment in women with polycystic ovary syndrome: A decision analysis of different oral ovulation induction agents. *Fertility and sterility* 2010;**94**(7):2659-64
23. Kar S. Clomiphene citrate or letrozole as first line ovulation induction drug in infertile PCOS women: A prospective randomised trial. *Fertility and sterility* 2012;**98**(3):S86
24. Kar S. Clomiphene citrate, metformin or the combination of both, as first line ovulation induction drug in polycystic ovarian syndrome: A randomised controlled trial. *Fertility and sterility* 2013;**100**(3):S359-S60
25. Ladson G, Dodson WC, Sweet SD, et al. The effects of metformin with lifestyle therapy in polycystic ovary syndrome: a randomized double-blind study. *Fertil Steril* 2011;**95**(3):1059-66 e1 doi: 10.1016/j.fertnstert.2010.12.002.
26. Legro RS, Brzyski RG, Diamond MP, et al. The Pregnancy in Polycystic Ovary Syndrome II study: baseline characteristics and effects of obesity from a multicenter randomized clinical trial. *Fertility & Sterility* 2014;**101**(1):258-69.e8 doi: <http://dx.doi.org/10.1016/j.fertnstert.2013.08.056>.
27. Legro RS, Brzyski RG, Diamond MP, et al. Letrozole versus clomiphene for infertility in the polycystic ovary syndrome. *Obstetrical and Gynecological Survey* 2015;**69**(10):599-601
28. Legro RS, Kunselman AR, Brzyski RG, et al. The Pregnancy in Polycystic Ovary Syndrome II (PPCOS II) trial: rationale and design of a double-blind randomized trial of clomiphene citrate and letrozole for the treatment of infertility in women with polycystic ovary syndrome. *Contemporary clinical trials* 2012;**33**(3):470-81 doi: 10.1016/j.cct.2011.12.005.


29. Legro RS, Myers ER, Barnhart HX, et al. The Pregnancy in Polycystic Ovary Syndrome Study: baseline characteristics of the randomized cohort including racial effects. *Fertility and Sterility* 2006;**86**(4):914-33 doi: <http://dx.doi.org/10.1016/j.fertnstert.2006.03.037>.
30. Lorzadeh N, Kazemirad S. Comparison of effects letrozole and clomiphene citrate for ovulation induction in women with polycystic ovary syndrome. *Journal fur Reproduktionsmedizin und Endokrinologie* 2010;**7**(4):367
31. Ma LK, Jin LN, Yu Q, et al. [Effect of lifestyle adjustment, metformin and rosiglitazone in polycystic ovary syndrome]. *Zhonghua Fu Chan Ke Za Zhi* 2007;**42**(5):294-7
32. Moll E, Korevaar JC, Bossuyt PMM, et al. Does adding metformin to clomifene citrate lead to higher pregnancy rates in a subset of women with polycystic ovary syndrome? *Human Reproduction* 2008;**23**(8):1830-34 doi: 10.1093/humrep/den182.
33. Moll E, van Wely M, Lambalk CB, et al. Health-related quality of life in women with newly diagnosed polycystic ovary syndrome randomized between clomifene citrate plus metformin or clomifene citrate plus placebo. *Human reproduction (Oxford, England)* 2012;**27**(11):3273-8 doi: 10.1093/humrep/des310.
34. Morin-Papunen L, Rantala AS, Unkila-Kallio L, et al. Metformin improves pregnancy and live-birth rates in women with polycystic ovary syndrome (PCOS): a multicenter, double-blind, placebo-controlled randomized trial. *Journal of Clinical Endocrinology & Metabolism* 2012;**97**(5):1492-500
35. Moslemizadeh N, Moghadam TG, Ehteshami S. Comparison of clomiphene citrate plus estradiol, with tamoxifen citrate effects in induction of ovulation and pregnancy in polycystic ovarian syndrome patients. *Journal of Medical Sciences* 2008;**8**(8):734-38
36. Nahid L, Sirous K. Comparison of the effects of letrozole and clomiphene citrate for ovulation induction in infertile women with polycystic ovary syndrome. [Italian]. *Minerva ginecologica* 2012;**64**(3):253-8
37. NIH/NICHD Reproductive Medicine Network. Letrozole versus clomiphene citrate in a novulatory PCOS women: A cost-effectiveness analysis. *Fertility and sterility* 2013;**100**(3):S128
38. NIH/NICHD Reproductive Medicine Network. Effect of Letrozole versus Clomiphene on live birth in women with anovulatory infertility due to polycystic ovary syndrome (PCOS): a randomized double-blind multi-center trial. *Fertility and sterility* 2013;**100** Suppl(3):S51
39. Rausch ME, Legro RS, Barnhart HX, et al. Predictors of pregnancy in women with polycystic ovary syndrome. *Journal of Clinical Endocrinology & Metabolism* 2009;**94**(9):3458-66 doi: 10.1210/jc.2009-0545.
40. Ridzuan J, Murizah MZ, Adibah I, et al. Prospective randomized control trial comparing metformin and clomiphene citrate as ovulation induction agent in women with polycystic ovarian syndrome at Alor Star hospital, Malaysia. XVIII FIGO World Congress of Gynecology and Obstetrics 2006;**3**:181-2
41. Schlaff D, Legro RS, Diamond MP, et al. Ovulation induction with clomiphene, metformin, or a combination of the two does not affect hirsutism score over a standard course of treatment in women with polycystic ovary syndrome. *Fertility and sterility* 2010;**94**(4):S192-S93

42. Seyedoshohadaei F, Zandvakily F, Shahgeibi SH. Effectiveness of clomiphene citrate, tamoxifen and letrozole in induction ovulation in women with infertility due to isolated unovulation. *Iranian Journal of Reproductive Medicine* 2011;**9**:33
43. Siebert TI, Kruger TF, Lombard C. Evaluating the equivalence of clomiphene citrate with and without metformin in ovulation induction in PCOS patients. *Journal of Assisted Reproduction & Genetics* 2009;**26**(4):165-71
44. Topcu HO, Yüksel B, İslimye M, et al. Tamoxifen as a first line treatment in ovulation induction for the infertile women with thin endometrium and high follicular response with clomiphene. *Human Reproduction* 2010;**25**:i312
45. Wang L, Lu S, Cao Z, et al. Effects of letrozole and clomiphene citrate on ovulation induction in polycystic ovarian syndrome. *Journal of Xi'an Jiaotong University (Medical Sciences)* 2008;**29**(6):69


Appendix 5 (a-e) Network plots of eligible comparisons for five outcomes: pregnancy, live birth, ovulation, miscarriage and multiple pregnancy.

The width of the lines is proportional to the number of trials comparing each pair of treatments, and the size of each node is proportional to the number of studies including the respective interventions. (Abbreviations: CC, clomiphene citrate; PB, placebo or no treatment; LET, letrozole; MF, metformin; TAM, tamoxifen; FSH, follicle stimulating hormone; LOD, laparoscopic ovarian drilling)


5a. Pregnancy




5b. live birth




5c. ovulation



5d. miscarriage




5e. multiple pregnancy



Appendix 6 Risk of bias evaluation.

6a. Risk of bias graph: review authors' judgments about each risk of bias item presented as percentages across all included studies.



6b. Risk of bias summary: review authors' judgments about each risk of bias item for each included study.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Abuelghar 2013	●	●	?	?	●	?	?
Amer 2009	●	●	●	●	●	●	●
Amer 2015	●	?	?	?	?	?	?
Atay 2006	?	?	?	?	●	?	?
Ayaz 2013	?	●	●	●	?	?	?
Aygen 2007	?	?	?	?	●	?	?
Badawy 2009	●	?	●	●	●	?	?
Badawy 2011	●	?	●	●	?	?	?
Basirat 2012	?	?	●	●	?	?	●
Bayar 2006	●	●	?	?	●	?	?
Beigi 2006	?	?	?	?	●	?	?
Boonstanfar 2001	●	●	●	●	●	?	?
Boudhraa 2010	?	?	?	?	?	●	?
Cudmore 1966	?	?	?	?	●	?	?
Dasari 2009	?	?	?	?	●	?	?
Dehbashi 2009	?	?	●	●	●	●	?
El-Biely 2001	●	?	?	?	?	?	?
Fleming 2002	●	●	●	●	●	?	●
Garcia 1985	?	?	●	?	●	?	?
Homburg 2012	●	●	●	●	●	●	●
Jahan 2015	?	?	?	?	?	?	?
Johnson 1966	?	?	?	?	●	?	?
Johnson 2010A	●	●	●	●	●	●	●
Johnson 2010B	●	●	●	●	●	●	●
Kar 2012	●	●	●	●	●	?	?
Kar 2013	●	●	●	●	●	?	?
Karimzadeh 2007	●	●	●	●	?	?	?
Karimzadeh 2010	?	?	?	?	?	?	?
Keikha 2011	●	●	●	●	?	?	?
Khorram 2006	●	?	●	●	?	?	?
Leanza 2014	●	●	●	●	●	?	?
Legro 2007	●	●	●	●	●	●	●
Legro 2014	●	●	●	●	●	●	●
Liu 2015	?	?	?	?	?	?	?
López 2004	●	●	?	?	●	?	?
Lord 2006	●	●	●	●	?	?	?
Lorzadeh 2011	?	?	?	?	●	?	?
Maged 2015	●	●	?	?	●	?	?
Mobusher 2014	?	?	?	?	●	?	●
Moll 2006	●	●	●	●	●	?	?
Nazik 2012	●	?	●	●	?	?	●
Palomba 2005	●	●	●	●	?	?	?
Raja 2005	?	?	?	●	●	?	?
Ray 2012	?	?	●	●	●	?	?
Robinson 2003	?	?	?	?	?	?	?
Roy 2012	●	●	?	●	●	?	?
Sahin 2004	?	?	?	?	●	?	?
Santonocito 2009	?	?	?	?	?	?	?
Selim 2012	●	●	●	●	●	?	?
Seyedoshohadaei 2012	?	?	●	?	●	?	?
Sharief 2015	?	?	?	?	●	?	?
Sh-EI-Arab Elseddek 2011	●	?	?	●	●	?	?
Tang 2006	●	●	●	●	●	?	?
Vegetti 1999	●	●	●	●	●	?	?
Williams 2009	?	?	?	?	?	?	?
Zain 2009	●	●	●	●	●	?	?
Zeinalzadeh 2010	?	?	?	?	●	?	?

Appendix 7 Pairwise meta-analysis results for direct comparisons of interventions

Comparisons		Pairwise meta-analysis odds ratio (95% CI)	No. of trials	No. of participants	Heterogeneity I ²
Pregnancy					
PB	vs CC	0.20(0.05-0.74)	3	136	0%
LET		1.53(1.26-1.85)	21	3553	24.3%
MF		1.10(0.62-1.95)	9	1335	73.1%
CC+MF		1.56(1.24-1.97)	19	2070	12.2%
TAM		0.64(0.36-1.12)	4	661	43.7%
FSH		1.57(1.04-2.37)	2	378	0%
LOD		0.52(0.19-1.44)	1	72	N/A
MF	vs PB	3.58(2.06-6.21)	5	494	0%
MF	vs LET	0.73(0.41-1.32)	1	304	N/A
TAM		0.67(0.30-1.47)	1	100	N/A
CC+MF	vs MF	1.92(0.90-4.06)	5	818	71.8%
Live birth					
LET	vs CC	1.60(1.30-1.98)	9	1990	0%
MF		1.00(0.45-2.22)	8	1155	80.9%
CC+MF		1.14(0.81-1.61)	7	950	12.4%
TAM		0.96(0.26-3.55)	2	195	35.3%
FSH		1.50(0.98-2.29)	2	378	0%
MF	vs PB	2.87(0.51-16.02)	1	65	N/A
MF	vs LET	0.38(0.19-0.78)	1	304	N/A
TAM		0.71(0.32-1.60)	1	100	N/A
CC+MF	vs MF	2.48(1.24-4.95)	4	640	51.1%
Ovulation (per woman randomised)					
PB	vs CC	0.15(0.07-0.34)	3	136	0%
LET		1.89(1.55-2.30)	14	2568	8.8%
MF		0.62(0.32-1.22)	7	1119	82.9%
CC+MF		1.46(1.01-2.12)	14	1407	54.5%
TAM		0.61(0.43-0.86)	3	566	0%
FSH		3.11(0.76-12.79)	1	76	N/A
LOD		0.70(0.27-1.83)	1	72	N/A
MF	vs PB	3.63(0.45-29.35)	3	309	92.9%
MF	vs LET	0.14(0.09-0.24)	1	304	N/A
TAM		0.75(0.31-1.78)	1	100	N/A
CC+MF	vs MF	3.20(1.85-5.52)	4	640	44.4%
Multiple pregnancy (per woman randomised)					
LET	vs CC	0.45(0.22-0.91)	12	2460	0%
MF		0.22(0.05-0.96)	4	976	0%
CC+MF		0.57(0.19-1.74)	4	892	0%
TAM		0.48(0.06-3.76)	2	471	0%


FSH		3.62(0.58-22.80)	2	378	0%
MF	vs PB	0.33(0.01-8.49)	1	65	N/A
MF	vs LET	0.20(0.01-4.15)	1	304	N/A
TAM		3.06(0.12-76.95)	1	100	N/A
CC+MF	vs MF	2.36(0.42-12.39)	4	665	0%
Miscarriage (per woman randomised)					
LET	vs CC	1.00(0.62-1.62)	10	2302	10.6%
MF		0.76(0.32-1.82)	8	1155	29.1%
CC+MF		1.38(0.85-2.24)	8	991	0%
TAM		0.56(0.19-1.68)	3	566	23.4%
FSH		1.44(0.57-3.63)	2	378	0%
MF	vs PB	1.02(0.28-3.73)	2	265	0%
MF	vs LET	0.33(0.13-8.20)	1	304	N/A
TAM		0.73(0.16-3.46)	1	100	N/A
CC+MF	vs MF	1.37(0.66-2.87)	4	640	10.9%
Miscarriage (per pregnant woman)					
LET	vs CC	0.79(0.52-1.21)	10	718	0%
MF		0.70(0.19-2.63)	8	277	54.9%
CC+MF		1.35(0.74-2.46)	8	384	0%
TAM		0.83(0.31-2.19)	3	123	0%
FSH		0.99(0.37-2.67)	2	164	0%
MF	vs PB	0.28(0.06-1.19)	2	63	0%
MF	vs LET	0.41(0.02-10.64)	1	55	N/A
TAM		0.93(0.18-4.72)	1	45	N/A
CC+MF	vs MF	0.67(0.27-1.66)	4	174	0%

(Abbreviations: CC, clomiphene citrate; PB, placebo or no treatment; LET, letrozole; MF, metformin; TAM, tamoxifen; FSH, follicle stimulating hormone; LOD, laparoscopic ovarian drilling)

An odds ratios < 1 favour the second intervention and an odds ratios > 1 favour the first intervention.

Appendix 8 Ranking of treatments for pregnancy


Rankograms below illustrate the probability per rank for each treatment in terms of pregnancy. E.g. for CC, the probabilities of being the best treatment, the second best, to the worst (eighth) are 0%, 0%, 2.4%, 29.0%, 55.5%, 12.3%, 0.8% and 0%, respectively.




Appendix 9 Network plot for pregnancy incorporating risk of bias assessment

9a. Risk of bias in randomisation

Coloured edges are based on adequacy of randomisation in the majority of the trials in each comparison. Green, yellow and red colours represent low, unclear and high risk, respectively.




9b. Risk of randomisation in allocation concealment




Appendix 10 Inconsistency plot for pregnancy

We estimated inconsistency as the logarithm of the ratio of two odds ratios (RoR) from direct and indirect evidence in the loop (also named inconsistency factor IF) and the corresponding 95% CI for each IF in each closed triangular or quadratic loop. RoR values is close to 1 mean that the two sources are in agreement. The inconsistency plot shows that in a total of 4 loops there is none with statistically significant inconsistency as all confidence intervals for RORs are compatible with zero inconsistency (RoR= 1).




Appendix 11 Comparison-adjusted funnel plot for pregnancy

The red line represents the null hypothesis that the study-specific effect sizes do not differ from the respective comparison-specific pooled effect estimates. Different colors correspond to different comparisons. (1-clomiphene; 2-placebo/no treatment; 3-letrozole; 4-metformin; 5-clomiphene plus metformin; 6-tamoxifen; 7-FSH; 8-LOD)




Appendix 12 Network meta-analysis results for live birth


The diamond in each line represents the estimate summary odds ratios of each comparison. The black lines represent the confidence intervals for summary odds ratios for each comparison and the red lines (overall length of the lines) the respective predictive intervals. The blue line is the line of no effect (odds ratio equal to 1). An odds ratio >1 favours the first intervention and an odds ratio < 1 favours the second. (Abbreviations: OR, odds ratio; CI, confidence interval; PrI, predictive interval; CC, clomiphene citrate; PB, placebo or no treatment; LET, letrozole; MF, metformin; TAM, tamoxifen; FSH, follicle stimulating hormone; LOD, laparoscopic ovarian drilling)




Appendix 13 Ranking of treatments for live birth




Appendix 14 Network meta-analysis results for ovulation




Appendix 15 Ranking of treatments for ovulation




Appendix 16 Network meta-analysis results for miscarriage per woman randomised




Appendix 17 Network meta-analysis results for miscarriage per pregnancy




Appendix 18 Network meta-analysis results for multiple pregnancy




Appendix 19 Ranking of treatments for multiple pregnancy



Appendix 20 Network plot for live birth incorporating risk of bias assessment
 20a. Risk of bias in randomisation




20b. Risk of bias in allocation concealment




Appendix 21 Inconsistency plot for live birth.


The inconsistency plot shows that in a total of 3 loops there is none with statistically significant inconsistency as all confidence intervals for RORs are compatible with zero inconsistency (RoR= 1).




Appendix 22 Comparison-adjusted funnel plot for live birth




Appendix 23 Network plot for ovulation incorporating risk of bias assessment
23a. Risk of bias in randomisation




23b.. Risk of bias in allocation concealment



Appendix 24 Inconsistency plot for ovulation




Appendix 25 Comparison-adjusted funnel plot for ovulation




(1-clomiphene; 2-placebo/no treatment; 3-letrozole; 4-metformin; 5-clomiphene plus metformin; 6-tamoxifen; 7-FSH; 8-LOD)


Appendix 26 Network plot for miscarriage incorporating risk of bias assessment
 26a. Risk of bias in randomisation




26b. Risk of bias in allocation concealment




Appendix 27 Ranking of treatments for miscarriage per pregnancy



Appendix 28 Inconsistency plot for miscarriage per pregnancy




Appendix 29 Comparison-adjusted funnel plot for miscarriage per pregnancy




(1-clomiphene; 2-placebo/no treatment; 3-letrozole; 4-metformin; 5-clomiphene plus metformin; 6-tamoxifen; 7-FSH)


Appendix 30 Network plot for multiple pregnancy incorporating risk of bias assessment
30a. Risk of bias in randomisation




30b. Risk of bias in allocation concealment



Appendix 31 Inconsistency plot for multiple pregnancy




Appendix 32 Comparison-adjusted funnel plot for multiple pregnancy




(1-clomiphene; 2-placebo/no treatment; 3-letrozole; 4-metformin; 5-clomiphene plus metformin; 6-tamoxifen; 7-FSH)


Appendix 33 Sensitivity analysis - RCTs reporting clinical pregnancy



Appendix 34 Sensitivity analysis - RCTs with treatment naïve women



Appendix 35 Sensitivity analysis - RCTs with low risk of randomisation & allocation bias



Appendix 36 Additional discussion

Side effects of the combination of clomiphene and metformin

We have summarised the side effects of the combination of clomiphene and metformin versus clomiphene alone (appendix 37 and 38). Of 19 studies comparing these two interventions, 11 studies reported data on side effects or discontinuation due to side effects. Three studies¹⁻³ including 714 women reported the number of participants who discontinued treatment due to side effects. In a pairwise meta-analysis for this outcome, we found that more women in the combination group discontinued the treatment due to side effects than women in clomiphene group (OR 2.34, 95% CI 1.04 to 5.30, appendix 37). As the reporting strategies were diverse in different studies, we were not able to perform meta-analyses on overall side effects or any specific types of side effects. As shown in appendix 38, gastrointestinal side effects were more frequent in combined clomiphene-metformin group than clomiphene group.

Quality of evidence and interpretation of data

The overall quality of included studies was moderate in relation to the seven specific domains of the risk of bias assessment. Randomisation and allocation are fundamental requirements for a high quality randomised controlled trial and therefore we integrated these domains in the network plot (appendix 9, 20, 23, 26, 30). Although we excluded quasi-randomised studies in the current systematic review, half of the included randomised controlled trials did not report details of randomisation, and further clarity on this eluded us even after attempts to contact the authors. Specific information about allocation concealment was also unavailable in many of the trials. In multicentre randomised controlled trials with large sample sizes^{1 2 4 5}, the dropout rates in different interventions varied from 14% to 35%. Many studies with small sample sizes have relatively low or zero dropout rates. Additionally, these studies often claim to have undertaken an intention-to-treat analysis, but it is possible that the authors may have excluded dropouts in their analysis. It is difficult to distinguish those lost to follow up due to adverse events and those for other reasons. CONSORT⁶ strongly encourages to report a flow diagram of patient follow up, including reasons for dropouts, however, many included studies failed to do so.

In pairwise meta-analyses, the heterogeneity in comparisons of combined clomiphene-metformin versus clomiphene and letrozole versus clomiphene in all outcomes was low. Therefore, the results of these comparisons in network meta-analysis were robust. By contrast, there was significant heterogeneity in comparisons of clomiphene and metformin. Thus, the results of these comparisons should be interpreted with cautions.

In our network meta-analysis, predictive intervals were used to estimate the effect of a future study. When considering predictive intervals in our network meta-analysis, clomiphene, letrozole, metformin, clomiphene and metformin combined, and follicle stimulating hormone remained superior to placebo. These results indicate that in future studies, these active treatments would remain effective in comparison with placebo/no treatment. Of note, there


were significant differences between follicle stimulating hormone and metformin/letrozole in terms of multiple pregnancy. However, the wide confidence intervals suggest significant imprecision in the effect size.

According to the rankings, combined clomiphene-metformin, letrozole, and follicle stimulating hormone were the best interventions in terms of pregnancy, live birth and ovulation, while metformin and letrozole were the best interventions in terms of reducing multiple pregnancy rate.

References

1. Legro RS, Barnhart HX, Schlaff WD, et al. Clomiphene, metformin, or both for infertility in the polycystic ovary syndrome. *New England journal of medicine* 2007;**356**(6):551-66.
2. Moll E, Bossuyt PM, Korevaar JC, et al. Effect of clomifene citrate plus metformin and clomifene citrate plus placebo on induction of ovulation in women with newly diagnosed polycystic ovary syndrome: randomised double blind clinical trial. *BMJ (Clinical research ed)* 2006;**332**(7556):1485.
3. Johnson NP, Stewart AW, Falkiner J, et al. PCOSMIC: a multi-centre randomized trial in women with PolyCystic Ovary Syndrome evaluating Metformin for Infertility with Clomiphene. *Human reproduction* 2010;**25**(7):1675-83.
4. Legro RS, Brzyski RG, Diamond MP, et al. Letrozole versus clomiphene for infertility in the polycystic ovary syndrome. *The New England journal of medicine* 2014;**371**(2):119-29.
5. Homburg R, Hendriks ML, Konig TE, et al. Clomifene citrate or low-dose FSH for the first-line treatment of infertile women with anovulation associated with polycystic ovary syndrome: a prospective randomized multinational study. *Human reproduction* 2012;**27**(2):468-73.
6. Moher D, Schulz KF, Altman D, et al. The CONSORT statement: revised recommendations for improving the quality of reports of parallel-group randomized trials. *Jama* 2001;**285**(15):1987-91.

Appendix 37 Meta-analysis of clomiphene and metformin combined vs clomiphene alone for discontinuation due to side effects.



Boxes and horizontal lines represent ORs and 95% CIs of individual studies. The diamond represents the overall OR and 95% CI (Random-effect model). OR >1 means more women discontinue treatment due to side effect in combined clomiphene-metformin group than clomiphene group.

Appendix 38 Side effects of clomiphene and metformin combined versus clomiphene alone.

Study ID	CC group				CC+Metformin group			
	Women with side effects	Details of side effects	discontinuation due to side effects	Sample size	Women with side effect	Details of side effect	discontinuation due to side effects	Sample size
Abuelghar 2013	6	Flushing: 4; gastrointestinal tract discomfort: 2	NA	32	11	Flushing: 2; gastrointestinal tract discomfort: 5; both: 1; diarrhoea: 3	NA	34
Ayaz 2013	NA	NA	0	21	NA	60% had complained of loss of appetite, 18% had nausea & vomiting	0	21
Basirat 2012	NA	No metformin related side effects.	NA	167	NA	No metformin related side effects.	NA	167
Dasari 2009	NA	NA	0	24	NA	NA ^a	0	16
Johnson 2010B	NA	Gastrointestinal symptoms: 5	1	36	NA	Gastrointestinal symptoms: 11; vasomotor: 1	0	35
Legro 2007	NA	Diarrhoea: 48; dyspepsia: 9; flatulence: 38; nausea: 82; stomach discomfort: 8; vomiting: 28; decreased appetite: 17 ^b	4	209	NA	Diarrhoea: 126; dyspepsia: 14; flatulence: 30; nausea: 138; stomach discomfort: 16; vomiting: 72; decreased appetite: 33 ^b	7	209
Maged 2015	1	Nausea: 1	NA	40	1	Drowsiness: 1	NA	40
Moll 2006	NA	NA	6	114	NA	NA	18	111
Raja 2005	NA	NA	NA	50	6	nausea and diarrhoea: 6	NA	50
Sahin 2004	NA	NA	0	10	NA	NA	0	11
Zain 2009	NA	NA	0	41	NA	NA ^c	0	41

NA: not available.

- a. The data of the 16 women in CC+ metformin group were not reported. But the authors reported that of the 25 participants who received metformin along with CC, 80% complained of loss of appetite and 24% had nausea and vomiting. The 25 participants was composed of 16 women in CC + metformin group and 9 women who did not conceive with six cycles of CC alone (given CC + metformin for an additional six cycles) at their request for further treatment.
- b. Main gastrointestinal side effects were summarised in this table. This study also reported data on other specific side effects but not the data on overall side effects.
- c. The data of CC+metformin group was not reported. Three patients with metformin complained of nausea, dizziness, and headache.

Appendix 39 Congenital malformations in newborns conceived through letrozole vs control.

Study ID	Country	Study design	Congenital malformation Control	Letrozole
Dehbashi 2009 ¹	Iran	RCT	CC: 16.6% (1/6) ^a	0% (0/10)
Ray 2012 ²	India	RCT	CC: 0% (0/13)	0% (0/20)
Roy 2012 ³	India	RCT	CC: 0% (0/21)	0% (0/39)
Legro 2014 ⁴	USA	RCT	CC: 1.5% (1/66) ^b	3.9% (4/102) ^c
Diamond 2015 ⁵	USA	RCT	CC: 4.3% (3/70) ^d	3.6% (2/56) ^e
Tulandi 2006 ⁶	Canada	observational	CC/CC+FSH: 4.8(19/397) ^f	Letrozole/Letrozole+FSH: 2.4% (14/514) ^g
Forman 2007 ⁷	Canada	observational	2.6% (7/271) ^h	0% (0/94)
Sharma 2014 ⁸	India	observational	CC:4.0% (10/251) ⁱ ; Natural conception: 2.9% (5/171) ^k	2.5% (5/201) ^j
Wu 2016 ⁹	China	RCT	Berberine: 0% (0/48)	Letrozole alone: 1.2%(1/84) ^l ; Letrozole+Berberine: 1.2%(1/81) ^m
Tatsumi 2016 ¹⁰	Japan	observational	Natural cycle IVF/ICSI: 1.9% (44/2287) ⁿ	Letrozole + IVF/ICSI: 2.2%(15/694) ^o

Details of congenital malformations in these studies:

a. Meningomyelocele.

b. Atrial septal defect (ASD), ventricular septal defect (VSD), and pulmonary stenosis.

c. 1) Cerebral palsy with arrested hydrocephalus with polycythemia and neutropenia; 2)

imperforate anus with perineal fistula and spina bifida with a tethered spinal cord; 3) right hemimegalencephaly, and dysgenesis of the left frontal and temporal lobes but no hydrocephalus; 4) large cardiac VSD requiring surgical repair.

d. 1) Aortic arch hypoplasia; 2) Congenital hypothyroidism; 3) Renal duplicated right collecting system and ureterocele.

e. 1) Hypospadias; 2) Right facial hemangioma; Biventricular hypertrophy; Bifid uvula; Small cataracts bilaterally; Widening of the corneal horizontal diameter.

- f. Major malformations (12 cases): 1) VSD (4 cases); 2) Transposition of great vessels; 3) Atresia of pulmonary valve and right ventricle; 4) Pulmonary valve atresia; 5) Pyelectasis; 6) Omphalocele; 7) Cleft palate; 8) Spinal muscular atrophy; 9) Down's syndrome.
 Minor malformations (7 cases): 1) Preauricular skin tag (2 cases); 2) Horseshoe kidney; 3) Polydactyly (3 cases); 4) Unspecific hypotonia.
- g. Major malformations (6 cases): 1) VSD; 2) Esophageal atresia; 3) Cleft palate; 4) Trisomy 18; 5) Down's syndrome; 6) Potter's syndrome.
 Minor malformations (8 cases): 1) Preauricular skin tag; 2) Congenital ptosis; 3) Plagiocephaly; 4) Hydrocele; 5) Hypospadias; 6) Polydactyly; 7) Syndactyly (2nd and 3rd toes); 8) Umbilical and inguinal hernias.
- h. 7 cases with major malformations but details not reported.
- i. 1) Patent ductus arteriosus (2 cases) and; 2) total anomalous venous connection; 3) Hypospadias (3 cases); 4) bilateral congenital talipes equino varus; 5) duplication of urethra; 6) cleft lip & palate; 7) inguinal hernia; 8) neural tube defect; 9) Down's syndrome (2 cases). Three babies with congenital heart disease were excluded from the analysis by the authors as they were born to diabetic mothers.
- j. 1) Combined ventricular and ASD; 2) paraumbilical hernia; 3) congenital deafness; 4) congenital talipes equino varus; 5) albinism.
- k. 1) VSD; 2) Congenital talipes equino varus; 3) cleft lip; 4) imperforate anus; 5) polydactyly.
- l. Hydrocephalus.
- m. Major VSD and pulmonary stenosis.
- n. Major anomalies (34 cases): including chromosomal abnormalities (11 cases), cardiovascular abnormalities (13 cases) and musculoskeletal abnormalities (1 case).
- o. Major anomalies (13 cases): 1) ASD, VSD; 2) ASD, VSD, Down's syndrome; 3) Cleft lip without cleft palate; 4) Congenital hydronephrosis; 5) Diaphragmatic hernia; 6) Duodenal atresia; 7) Endocardial cushion defect, down syndrome; 8) Hypospadias; 9) Trisomy 18; 10) VSD (2 cases); 11) VSD, down syndrome; 12) Anencephalus.

References:

1. Dehbashi S, Dehbashi S, Kazerooni T, et al. Comparison of the effects of letrozole and clomiphene citrate on ovulation and pregnancy rate in patients with polycystic ovary syndrome. *Iranian Journal of Medical Sciences* 2009;**34**(1):23-28.
2. Ray PB, Ray A, Chakraborti PS. Comparison of efficacy of letrozole and clomiphene citrate in ovulation induction in Indian women with polycystic ovarian syndrome. *Archives of Gynecology & Obstetrics* 2012;**285**(3):873-7.
3. Roy K, Baruah J, Singla S, et al. A prospective randomized trial comparing the efficacy of Letrozole and Clomiphene citrate in induction of ovulation in

polycystic ovarian syndrome. Journal of human reproductive sciences 2012;**5**(1):20-25.

4. Legro RS, Brzyski RG, Diamond MP, et al. Letrozole versus clomiphene for infertility in the polycystic ovary syndrome. The New England journal of medicine 2014;**371**(2):119-29.
5. Diamond MP, Legro RS, Coutifaris C, et al. Letrozole, Gonadotropin, or Clomiphene for Unexplained Infertility. The New England journal of medicine 2015;**373**(13):1230-40.
6. Tulandi T, Martin J, Al-Fadhli R, et al. Congenital malformations among 911 newborns conceived after infertility treatment with letrozole or clomiphene citrate. Fertility and sterility 2006;**85**(6):1761-5.
7. Forman R, Gill S, Moretti M, et al. Fetal safety of letrozole and clomiphene citrate for ovulation induction. J Obstet Gynaecol Can 2007;**29**(8):668-71.
8. Sharma S, Ghosh S, Singh S, et al. Congenital malformations among babies born following letrozole or clomiphene for infertility treatment. PloS one 2014;**9**(10):e108219.
9. Wu XK, Wang YY, Liu JP, et al. Randomized controlled trial of letrozole, berberine, or a combination for infertility in the polycystic ovary syndrome. Fertility and sterility 2016;**106**(3):757-65 e1.
10. Tatsumi T, Jwa SC, Kuwahara A, et al. No increased risk of major congenital anomalies or adverse pregnancy or neonatal outcomes following letrozole use in assisted reproductive technology. Human reproduction 2016.